

*Atlantic Sunrise Project – PA DEP Chapter 105 Joint Permit Application  
Transcontinental Gas Pipe Line Company, LLC  
Wyoming County*

**ATTACHMENT L -4  
ENVIRONMENTAL ASSESSMENT FORM ENCLOSURE C**

**ENCLOSURE C  
DESCRIPTION OF AQUATIC HABITAT**

Transcontinental Gas Pipe Line Company, LLC (Transco) is submitting an application to the Pennsylvania Department of Environmental Protection (PA DEP) for a Joint Permit under the Chapter 105 Pennsylvania Water Obstruction and Encroachment guidelines. This permit application only addresses the portions of the Project located in Pennsylvania, subject to PA Code Title 25 Chapter 105 and within the jurisdiction of the Northeast, North Central and South Central Regions of the PA DEP. For Project-related impacts to waters of the United States subject to jurisdiction under Section 404 of the Clean Water Act, Transco has applied for an Individual Permit with the U.S. Army Corps of Engineers (USACE) Baltimore District. Transco applied for a 401 Water Quality Certification for the entire Project on April 9, 2015 at the direction of PA DEP. Attachment J of this permit application includes a Project Description that further details the purpose and need and components of the proposed Atlantic Sunrise Project.

The Project is located primarily within the Susquehanna River drainage basin subregion (4-digit HUC) which is within the Upper Susquehanna Major Basin (6-digit HUC) and the Upper Susquehanna-Tunkhannock Subbasin (8-digit HUC). The Project is located in watersheds with water quality classifications identified as Cold Water Fishes (CWF) and Migratory Fishes (MF), Warm Water Fishes (WWF) and MF, and Trout Stocked Fishes (TSF) and MF according to PA Code Title 25 Chapter 93.

The Wetland Delineation and Stream Identification Report in Section L, Enclosure A, provides additional information regarding specific streams and waterbodies in the Project study area. The total construction workspace associated with the Project located in Wyoming County, Pennsylvania is approximately ~~378.51~~ **440.79** acres. To date, 100 percent of the workspace (22.5 pipeline miles) within Wyoming County has been surveyed for aquatic habitats including wetlands and waterways. ~~Only those resources that have been identified in the field (ground truthed) are included in this Environmental Assessment Form.~~ The *surveyed* wetland and waterbody boundaries ~~for ground truthed resources were~~ *were initially* reviewed in the field and verified by the USACE and PA DEP in June 2015. **Additional field reviews with USACE and PA DEP were conducted in August 2016, December 2016, and April 2017.**

Field surveys conducted to date identified **49-59** wetlands and **46-51** watercourses located within the Project construction workspace in Wyoming County.

## **A. Aquatic Habitat**

### **A1. Food Chain Production**

Intermittent and perennial streams in the Project study area contribute to aquatic and terrestrial food chains by serving as a breeding site for fish, insects, and amphibians. Larger wetlands that are inundated for extended periods of time may also contribute to food chain production. The larval forms of many insects and macroinvertebrates are aquatic, and amphibians spend all or part of their life cycle in aquatic habitats. Insects and amphibians produced in aquatic habitats can be locally important food organisms for a variety of fish, reptiles, birds, and mammals. Moreover, wetlands include plant food sources that may be used by a number of bird, and mammal species. Emergent plant species providing food value to wildlife include bulrush, sedge, and grass species. Shrub and woody vegetation growing within wetlands identified as PFO and PSS contain plants that may be used as a food source by various bird species. PEM wetlands can provide food to various bird species as well.

### **A2. General Habitat**

Transco delineated and classified wetland resources within the Project area through field surveys conducted in 2014, 2015, ~~and 2016~~, **and 2017**. Wetlands crossed by the Project in Pennsylvania were field delineated in accordance with the United States Army Corps of Engineers USACE Wetlands Delineation Manual (USACE 1987), **the Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0) (USACE 2011)**.

The USFWS wetland classification system described by Cowardin, et al. (1979) was used to classify wetlands delineated within the Project area. The wetlands within the Project area were identified as palustrine forested (PFO), palustrine scrub-shrub (PSS), and palustrine emergent (PEM), or a combination of these three Cowardin classes. Palustrine systems include all non-tidal wetlands that are dominated by trees, shrubs, persistent emergent, and emergent mosses or lichens and all wetlands that occur in tidal areas where salinity due to ocean-derived salts is

below 0.5 percent. The palustrine system was developed to group vegetated wetlands, commonly referred to as marshes, swamps, bogs, and prairies. This system includes ponds and may be situated shoreward of lakes, river channels, estuaries, and river floodplains or in isolated catchments or on slopes (Cowardin et al. 1979).

To obtain regional relevance, the Cowardin classes can be described according to the *Terrestrial and Palustrine Plant Communities of Pennsylvania* (Fike 1999). Vascular plant species nomenclature presented in this section is from *The Plants of Pennsylvania* (Block and Rhoads 2007). The classification system used by Fike (1999) defines plant communities as “an assemblage of plant populations sharing a common environment and interacting with each other, with animal populations, and with the physical environment.” Though Fike classifications were not specifically conducted for each delineated wetland, those described in this chapter were noted to be present and associated with at least some of the mapped wetlands within the Project area.

#### Palustrine Forested Wetlands (PFO)

Cowardin class PFOs are characterized by woody vegetation that is 6 meters (approximately 20 feet) tall or taller and normally include an overstory of trees, an understory of young trees or shrubs, and an herbaceous layer (Cowardin et al. 1979). According to *Terrestrial and Palustrine Plant Communities of Pennsylvania* (Fike 1999), areas in which the canopy is closed or nearly closed, with 60 to 100 percent tree cover, are classified as Palustrine Forests. Areas in which the tree cover is sparse (between 10 and 60 percent cover) are classified as Palustrine Woodlands. Shrubs may be present in these areas, but they do not dominate the community. Herbaceous and non-vascular plant covers also are known to occur in Palustrine Woodlands. Fike (1999) palustrine forest communities correlating to Project area PFO wetlands include the following:

##### *Coniferous Palustrine Forests: Hemlock - Mixed Hardwood Palustrine Forest*

This community type is dominated by a mixture of conifers and hardwood species and typically is characterized by a mineral soil or muck over mineral soil and by some groundwater enrichment. Eastern hemlock (*Tsuga canadensis*) comprises between 25 and 75 percent of the canopy. Other conifer species include white pine (*Pinus strobus*) and red spruce (*Picea rubens*). Yellow birch (*Betula lenta*), red maple (*Acer rubrum*), black ash (*Fraxinus nigra*), black gum

(*Nyssa sylvatica*), and gray birch (*Betula populifolia*) are the predominant hardwood species found in these communities. Common shrubs include highbush blueberry (*Vaccinium corymbosum*), winterberry (*Ilex verticillata*), and swamp azalea (*Rhododendron viscosum*). Typical herbaceous vegetation includes cinnamon fern (*Osmunda cinnamomea*), sensitive fern (*Onoclea sensibilis*), sedge species (*Carex* spp.), and skunk cabbage (*Symplocarpus foetidus*). Sphagnum moss (*Sphagnum* spp.) often is found in hemlock-mixed hardwood palustrine forest communities (Fike 1999).

*Coniferous - Broadleaf Palustrine Woodlands: Hemlock - Mixed Hardwood Palustrine Woodland*

This category describes a group of wetland forests dominated by a mixture of conifers and hardwood species. The substrate usually is mineral soil or muck over mineral soil. There is generally some groundwater enrichment in these systems. Eastern hemlock contributes 25 percent to 75 percent of the canopy. Other conifer species that may occur with hemlock include eastern white pine, red spruce, and tamarack (*Larix laricina*). The most common hardwood species are yellow birch, red maple, black ash, black gum, and gray birch. Rosebay (*Rhododendron maximum*) often forms a dense understory; other shrubs include highbush blueberry, winterberry, swamp azalea, and witherod (*Viburnum cassinoides*). Herbaceous species include cinnamon fern, sedges (e.g., *Carex trisperma*, *Carex prasina*, *Carex leptalea*), violets (*Viola* spp.), skunk-cabbage, false hellebore (*Veratrum viride*), sensitive fern, partridge-berry (*Mitchella repens*), gold-thread (*Coptis trifolia*), Canada mayflower (*Maianthemum canadense*), crested wood fern (*Dryopteris cristata*), and purple-stemmed aster (*Symphotrichum puniceum*). The bryophyte layer usually is well developed and may be dominated by sphagnum (Fike 1999).

*Broadleaf Woodlands: Red Maple – Mixed Shrub Palustrine Woodland*

This community type is characterized by a tree cover between 10 and 60 percent, dominated by red maple, and has a dense understory of shrubs. The substrate usually is mineral soil or muck over mineral soil. Other tree species may include blackgum (*Nyssa sylvatica*), Eastern hemlock, eastern white pine, black willow (*Salix nigra*), swamp white oak (*Quercus bicolor*), pin oak (*Q. palustris*), and black ash. The shrub layer is typically dense and includes silky dogwood (*Cornus amomum*), winterberry, spicebush (*Lindera benzoin*), smooth alder (*Alnus serrulata*), silky willow (*Salix sericea*), swamp rose (*Rosa palustris*), and buttonbush (*Cephalanthus occidentalis*). Ferns typically are the dominant vegetation within the herbaceous layer and include marsh fern

(*Thelypteris palustris*), cinnamon fern, sensitive fern, crested wood fern (*Dryopteris cristata*), and royal fern (*Osmunda regalis*). Other herbs include skunk-cabbage, beggar-ticks (*Bidens* spp.), and jewelweed (*Impatiens capensis*). Areas of the woodland floor with long-duration hydrology support the growth of arrow-arum (*Peltandra virginica*), wapato (*Sagittaria latifolia*), and marsh-marigold (*Caltha palustris*) (Fike 1999).

#### Palustrine Scrub-Shrub Wetlands (PSS)

Cowardin class PSS wetlands include freshwater wetlands dominated by woody vegetation less than 20 feet in height or with trunks less than 3 inches diameter at breast height (dbh). PSS wetlands include true shrubs, saplings, young trees, and trees or shrubs that are small or stunted because of environmental conditions (Cowardin et al. 1979). PSS wetlands are widespread, highly variable communities with shrub-dominated wetlands that occur on mineral or mucky mineral soils that are either seasonally or temporarily flooded. They are typically found in flat areas in which the water table is at or above the soil surface for most of the year. Shrub swamps generally are found on the transition zone of emergent and forested areas that have been previously disturbed.

Review of Project area PSS wetlands did not identify any direct correlations between this Cowardin class and Fike (1999) palustrine shrubland communities. Regardless, typical vegetation within Project area PSS wetlands include multiflora rose (*Rosa multiflora*), black willow, red maple, highbush blueberry, tussock sedge (*Carex stricta*), and jewelweed. Multiflora rose is listed as a noxious weed within the Commonwealth of Pennsylvania.

#### Palustrine Emergent Wetlands (PEM)

Cowardin class PEM wetlands are non-tidal wetlands characterized by erect, rooted, herbaceous hydrophytes, excluding mosses and lichens. Hydrophytic vegetation is present for a majority of the growing season in most years. PEM wetlands usually are dominated by perennial herbaceous plants, though some woody plants may be present. Plant communities present in PEM wetlands can be very dynamic due to the variable hydrologic conditions through the growing season and between growing seasons (Cowardin et al. 1979). Fike (1999) herbaceous emergent wetland communities correlating to Project area PEM wetlands include the following:

*Emergent Wetland: Mixed Forb – Graminoid Wet Meadow*

This is a common, statewide, diverse wetland type characterized by poorly drained clay loam soils, few to scattered trees, and may occur as old fields, grasslands, or alluvial bottoms where trees have been removed. This type of wetland may be subject to seasonal grazing or mowing. Plant species composition is diverse, though some sites may be dominated by one or two species. Representative species include a combination of grass, grass-like (Graminoid) and forb species. Species include goldenrods (*Solidago* spp.), rice cutgrass (*Leersia oryzoides*), woolgrass (*Scirpus cyperinus*), bugleweed (*Lycopus uniflorus*), smartweeds, sedges (*Carex stipata* var. *stipata*, *C. canescens*, *C. lurida*, *C. cristatella*, *C. tribuloides*, *C. vesicaria*, *C. stricta*), soft rush (*Juncus effusus*), Joe-Pye-weed (*Eutrochium* spp.), boneset (*Eupatorium perfoliatum*), cinnamon ferns, Canadian St. John's-wort (*Hypericum canadense*), bluejoint (*Calamagrostis canadensis* var. *canadensis*), New York ironweed (*Vernonia noveboracensis*), beggar-ticks (*Bidens* spp.), dwarf St. John's-wort (*Hypericum mutilum*), bulrush (*Scirpus* spp.), marsh St. John's-wort (*Triadenum virginicum*), rattlesnake mannagrass (*Glyceria canadensis*), and spike-rushes (*Eleocharis* spp.). Scattered shrubs may be present, including steeplebush (*Spiraea tomentosa*), silky dogwood (*Cornus amomum*), gray dogwood (*Cornus racemosa*), red-osier dogwood (*Cornus sericea*), and arrow-wood (*Viburnum recognitum*). Noxious weed species such as purple loosestrife (*Lythrum salicaria*) and a variety of non-native grasses, such as reed canary-grass (*Phalaris arundinacea*) are frequently found in these meadows (Fike 1999)

*Emergent Wetland: Mixed Forb Marsh*

This wetland community occurs along lake margins, flooded depressions, and other wetlands that remain inundated throughout the growing season. Soils may contain muck at or near the surface and many of these wetlands are associated with sandy soils. The plant composition is variable and includes aquatic emergent plants as well as submerged aquatic species. Characteristic species include three-way sedge (*Dulichium arundinaceum* var. *arundinaceum*), halberd-leaf tearthumb (*Persicaria arifolia*), tearthumb (*Persicaria sagittata*), dock (*Rumex* spp.), sharp-fruited rush (*Juncus acuminatus*), beggar-ticks (*Bidens* spp.), jewelweed, sensitive fern, wapato, sedges (*Carex stricta*, *C. lacustris*, *C. lurida*, *C. crinita*, *C. stipata*, *C. tribuloides*, *C. scoparia*, *C. projecta*, *C. comosa*, *C. hystericina*, *C. baileyi*), sweet-flag (*Acorus calamus*), and rice cutgrass (Fike 1999).

**Emergent Wetland: Skunk Cabbage – Golden Saxifrage Seep**

Emergent wetlands are associated with seeps that occur where groundwater flow is expressed at the surface and saturates a broad area without forming a channel, which would otherwise promote drainage. These wetland types often form underneath a forest canopy. Species composition is highly variable, but can include turtlehead (*Chelone glabra*), red maple, Jack-in-the-pulpit (*Arisaema triphyllum*) Pennsylvania bittercress (*Cardamine pensylvanica*), sedge (*Carex prasina*), winterberry, spicebush, hornbeam (*Carpinus caroliniana*), skunk-cabbage, golden saxifrage (*Chrysosplenium americanum*), cinnamon fern, sedge (*Carex scabrata*), jewelweed, spinulose wood fern (*Dryopteris carthusiana*), Pennsylvania bittercress (*Cardamine pensylvanica*), clearweed (*Pilea pumila*), slender mannagrass (*Glyceria melicaria*), swamp saxifrage (*Saxifraga pensylvanica*), and sensitive fern. Bryophytes include *Brachythecium rivulare*, *Bryhnia novae-angliae*, *Rhynchostegium serrulatum*, *Rhizomnium punctatum*, *Sphagnum* spp., and *Thuidium delicatulum*. Shrubs may be present, common representatives include alders (*Alnus* spp.), spicebush, viburnums (*Viburnum* spp.), and dogwoods (*Cornus* spp.) (Fike 1999).

Information on aquatic habitats was derived from three principal sources: Project-specific communication with federal and state agencies; published and unpublished natural resources data pertaining to the regional Project area; and biological field surveys. The field surveys included wetland delineations and rare species habitat evaluations. Table L-4(c)-1 lists the ecological functions served by each wetland and waterbody type within Wyoming County.

**Table L-4(c)-1  
 Ecological Functions of Wetlands Associated with the Atlantic Sunrise Project within Wyoming County**

Ecological Function	Emergent Wetlands	Forested Wetlands	Scrub-Shrub Wetlands
Groundwater Re-charge/Discharge	X		
Floodflow Alteration	X	X	X
Fish and Shellfish Habitat	X	X	X
Sediment/Toxicant/Pathogen Retention	X	X	
Nutrient Removal/Retention/Transformation	X		
Production/Export	X		
Sediment/Shoreline Stabilization		X	
Wildlife Habitat	X	X	X
Recreation			

**Table L-4(c)-1**  
**Ecological Functions of Wetlands Associated with the Atlantic Sunrise Project within Wyoming County**

Ecological Function	Emergent Wetlands	Forested Wetlands	Scrub-Shrub Wetlands
Education/Scientific Value			
Uniqueness/Heritage	X	X	X
Visual Quality/Aesthetics			
Threatened or Endangered Species Habitat			

Riverine

Riverine systems are defined as wetland and deepwater habitats contained within a channel with periodic or continuously moving water. Although a multitude of intermittent and perennial stream systems are crossed by the Project route, the Susquehanna River is the most prominent Riverine feature represented within the Project area. According to Fike (1999) sparsely vegetated wetland communities, two communities have the potential to be present along the Susquehanna River: Floodplain Scour Community and Periodically Exposed Shoreline Community. Review of the crossing locations along the Project route did not identify the presence of either of these communities.

Sparsely Vegetated Wetland Communities - Periodically Exposed Shoreline Community

This community is found on all orders of streams, and in a wide variety of riverine settings including island heads, bars, spits, low terraces, and river banks. The underlying substrate also varies greatly, though it is often cobbles and sand with thin deposits of silt, muck, or organic matter.

According to Fike (1999), this community type is a component of ponds, lakes, and river floodplains, and occurs along nearly every creek and river. This community type, while common, provides important habitat for a number of rare insect species, namely tiger beetles (genus *Cicindela*), and ground beetles (order *Carabidae*) and others.

Aside from the Fike community described above, rivers/streams within the Project route have many values assigned to them by the state of Pennsylvania. Based on Pennsylvania Code (Title 25, Chapter 93), streams designated as Trout Stocking Fisheries have conditions suitable for supporting stocked trout between February 15 and July 31, and maintaining other flora and

fauna indigenous to a warm water habitat. High Quality Waters support a high quality benthic macroinvertebrate and fish community, potentially including trout. Exceptional Value Waters are similar to High Quality Waters, but may maintain a healthier macroinvertebrate community and/or be located in a designated area that supports natural communities such as a state park.

Under Pennsylvania Code Title 58, Chapter 57, the Pennsylvania Fish and Boat Commission (PFBC) classifies waterbodies that support trout populations (either stocked or native) or provide trout habitat as follows: Wild Trout Waters (also identified by PFBC as “Wild Trout Streams”, “Stream Sections that Support Natural Reproduction of Trout” and “Trout Natural Reproduction Waters”), including upstream tributaries; Class A Wild Trout Streams; and Wilderness Trout Streams. In general, Wilderness Trout Streams provide the most suitable conditions for trout as reflected by natural abundance levels, followed by Class A Wild Trout Waters and then Approved Trout Waters. PFBC-designated Trout Stocked Streams and Warmwater/Coolwater fisheries are generally not cold enough to maintain trout populations, but provide suitable habitat for seasonal stocking of trout and other sport fish.

#### **A2a through A2g. Nesting, Spawning, Rearing, Resting, Migration, Feeding, and Escape Cover**

Within the Project route, these forested wetland communities provide habitat for a variety of avian, mammal and amphibian species. For avian species, vegetation structure provides nesting, rearing, resting, escape cover, feeding, and migratory habitat. The intermittent water table, which excludes the presence of fish populations within these wetlands, provides diverse habitat for amphibians, including spawning and feeding habitats. Amphibian habitat is often provided by a “hummock and hollow” microtopography within this palustrine forest community (Fike 1999). Field delineated wetlands within this forest type for the Project noted that “hummock and hollow” micro-topography was often present where Eastern hemlocks (*Tsuga canadensis*) occupy hummocks, and hollows occur between trees.

South Branch Tunkhannock Creek and ~~Martin~~ **Tunkhannock** Creek are classified by the PA Fish and Boat Commission (PFBC) as approved trout waters. The main stem of the South Branch of Tunkhannock Creek has been confirmed by PFBC to be stocked within 0.5 mile of the Project area. Additionally, **Leonard Creek, Marsh Creek, Mill Creek, Mill Run, and Whitelock Creek, and their tributaries** is are classified by PFBC under PA Code Title 58, Chapter 57.11

as a Wild Trout Stream, or a stream that supports the natural reproduction of trout.

**Additionally, three UNTs to Leonard Creek are designated by PFBC as Class A Wild Trout Stream within the Project area.** The Susquehanna River is also listed by PFBC as a Warmwater/Coolwater Fishery (WWCW). The perennial waterbodies in the Project study area may provide spawning habitat for trout and/or small fish species and contribute to the water quality of downstream waters for the spawning of larger fish. The substrate within each of the waterbodies crossed by the Project is variable and includes boulder, cobble, gravel, sand, silt, clay, and vegetation. All waterbodies in the Project area are considered MF for the passage, maintenance, and propagation of anadromous and catadromous fishes and other fishes which move to, or from, flowing waters to complete their life cycle in other waters, according to PA Code Title 25 Chapter 93. It has been confirmed with PFBC that there are currently no stocked anadromous fishes within the Project area. Adequate fish cover is available under cobble, undercut banks, and woody debris.

A variety of open water and vernal pool aquatic habitats were delineated within the Project route. Some of these may be classified as vernal pools, but wetland survey methods did not specifically target the identification of vernal pools. Due to long duration ponded water, these aquatic resources are typically sparsely vegetated, and in the case of vernal pools, susceptible to seasonal drying.

Waterbodies within the Project area contain cobble and woody debris that may provide a resting habitat for aquatic organisms. Vegetation within wetlands provides shade and limited resting opportunities for wildlife species such as small mammals, amphibians, and insects. Larger PEM, PSS, and PFO wetlands with a greater degree of vegetation profile heterogeneity may provide additional resting habitat and escape cover for wildlife species.

Riffle-pool complexes in the streams provide minimal escape cover for aquatic species. Woody debris and undercut banks may also serve as cover for escape. Some wetlands in the Project area have sufficient vegetation to provide escape cover for small vertebrates and white-tailed deer.

Since PA falls within both the Atlantic and Appalachian Flyways for migratory birds, many bird groups utilize PA's natural habitats for resting, breeding, and nesting. Breeding populations of

interior forest bird species depend upon unfragmented forests. **No Important Bird Areas (IBAs) are crossed by the Project within Wyoming County.** Transco developed a Migratory Bird Plan in coordination with the USFWS Pennsylvania Field Office. The plan outlines conservation measures proposed to minimize potential impacts on migratory birds. ***Transco's Migratory Bird Plan includes compensatory mitigation to offset the removal of upland forest and forest fragmentation, and was approved by USFWS on February 6, 2017. Transco is also developing a Memorandum of Agreement (MOA) with the USFWS that includes mitigation to offset the removal of upland forest and forest fragmentation.***

#### **A2h. Other**

**No other special aquatic habitats are crossed by the Project in Wyoming County.**

#### **A3. Habitat for Threatened and Endangered Plant and Animal Species**

Coordination has been initiated with the PA DCNR, PFBC, PGC and USFWS, and field surveys have been completed to determine the presence of sensitive plant and wildlife species within the Project area. A summary of this correspondence and field survey results is included in Attachment G of this permit application package.

#### **A4. Environmental Study Areas**

The Project is not located near any known Environmental Study Areas.

##### **A4a. Sanctuaries**

The Project will not affect areas dedicated for use as sanctuaries by state or federal agencies or non-profit organizations.

##### **A4b. Refuges**

The Project will not affect areas dedicated for use as refuges.

#### **A5. Description of Instream Macroinvertebrate Communities**

The Project does not involve stream relocation, placement of fill, placement of a water obstruction or dredging. cursory visual inspections were recorded for streams in the Project area, which will be temporarily impacted by construction. Macroinvertebrates and amphibians

observed were recorded in Section L, Enclosure A on each waterbody's corresponding data sheet.

## **B. Water Quantity and Streamflow**

### **B1. Natural Drainage Patterns**

The Upper Susquehanna River Basin drains 424 square miles in Pennsylvania, with largest contributing waterbody crossed by the Project within Wyoming County being the Susquehanna River. Post-construction contours will be returned to approximately the same as pre-construction contours, leaving existing drainage patterns intact. Trench plugs will be installed at all resource crossings to maintain existing hydrology within wetlands and waterbodies. Crossing methods for waterbodies have been designed to accommodate the drainage area and flow characteristics of each particular crossing.

### **B2. Flushing Characteristics**

Large storm events and spring snow melt likely flush the streams and wetlands in the area. Some wetlands drain to unnamed secondary channels and may contribute flushing flows to streams within the watershed.

### **B3. Current Patterns**

Streams in the Project area do not affect current patterns in other surface waters downstream of the Project area. Wildlife activity (i.e., beaver dams) may alter water flow out of wetland areas into waterbodies.

### **B4 and B5. Groundwater Discharge for Baseflow Natural Recharge Area for Ground and Surface Waters**

The Valley and Ridge aquifer is located underneath the Project for its entirety. The aquifer is 0 to 155 feet below ground surface and has an average well yield of 2 to 215 gallons per minute. Precipitation, adjacent underlying bedrock, and stream flow recharge these aquifers. Most recharge of the glacial aquifer is discharged to the Susquehanna River or withdrawn throughout the watershed via potable or non-potable water supplies. Wetlands in the Project area may provide groundwater or surface water recharge, depending on soil permeability.

## **B6. Storm and Flood Water Storage and Control**

Streams and wetlands in the Project area may function to attenuate flood waters and provide flood storage or control. During storm events, the Project area wetlands store stormwater that may drain to large perennial waterbodies. Wetlands within the Project area would provide some capacity to store flood waters and may serve to reduce the severity of flood peaks emanating from their upstream watersheds.

## **C. Water Quality**

### **C1. Preventing Pollution**

Pollution prevention is an aquatic function that includes a variety of specific pollutant removal mechanisms such as trapping sediments, removal and burial of metals and other toxic compounds, and biological transformation and degradation of nutrients and organic pollutants. Some of these pollutants may become trapped in the substrate and deposit within streams and wetlands in the Project area. A wetland's ability to improve water quality depends on the type and density of aquatic vegetation, reduction and oxidation (redox) conditions and organic content of the sediments, available water storage volume, and detention time within the wetland. Most of the wetlands also have a depositional environment, which aids in trapping and retaining pollutants.

Transco obtained federal and state search reports from Environmental Data Resources, Inc. (EDR) to determine the presence and location of potential groundwater contamination in the vicinity of all proposed pipeline facilities and new aboveground facilities. The search area was based on a 2-mile radius extending from the centerline of the pipeline loop to account for any subsequent route modifications. **No sites with potential groundwater contamination were identified within the search areas for the Project in Wyoming County.**

For proposed modifications to existing aboveground facilities that are not within the search radius for the pipeline EDR reports, Transco completed a search for potential sources of contaminated groundwater using the USEPA's multisystem database. The search of the USEPA's multisystem database for records of potential groundwater contamination within 0.25 mile of all existing compressor stations, meter/regulator stations, contractor/pipe yards, and

MLVs did not identify any potential groundwater contamination in the vicinity of the aboveground facilities associated with the Project in **Wyoming County**.

## **C2. Sedimentation Control and Patterns**

Sediment stabilization includes stream bank anchoring, dissipation of erosion forces and trapping of sediments. Siltation observed within some of the Project area streams demonstrated in-stream sedimentation. In general, any vegetated cover (e.g., herbaceous and scrub/shrub vegetation) within wetlands can stabilize soils and trap sediments. Some portions of the wetlands in the Project area contain dense vegetation that aids in retaining sediments and filtering water.

## **C3. Salinity Distribution**

The Project is not located in a coastal or estuarine environment, therefore, this section is not applicable.

## **C4. Natural Water Filtration**

Streams in the Project area may perform some functions in water filtration; there is moderate variation in the stream substrate and/or gradient. Streambed substrate observed in streams most commonly included silt, sand, gravel, and cobble, but also included vegetation, boulder, and bedrock. Wetlands, especially those with dense vegetation, can effectively detain and naturally filter water laden with nutrients and/or toxicants. In the Project area, the wetlands and riparian vegetation abutting or adjacent to streams serve to some extent to maintain natural water filtration. Most emergent portions of wetlands within the Project area contain areas of dense vegetation that could aid in the natural water filtration process.

## **D. Recreation**

### **D1 and D2. Game and Non-Game Species**

The Atlantic Sunrise Project area traverses habitats that support a variety of wildlife species. Vegetative cover type and density are important environmental factors influencing wildlife habitat and species distribution. Variations in vegetative community types (e.g., deciduous hardwood and conifer are community types within the forested upland vegetation cover type) and other conditions, such as topography and existing land use, influence the quality and

availability of wildlife habitat within the Project area. Areas of greatest wildlife diversity and density are expected in natural habitats such as extensive, contiguous forest tracts, successional habitats (scrub-shrub), and grasslands. Habitats in agricultural land such as pastures, croplands, and hayfields harbor generalist wildlife species, consisting primarily of small mammals and white-tailed deer.

### Upland Forest

Upland forests provide food resources, cover, and nesting habitat for a variety of fauna in the vicinity of the Project. The tree canopy and shrub understory provide food and cover for larger mammals such as white-tailed deer (*Odocoileus virginianus*) and black bear (*Ursus americanus*). Leaf litter on the forest floor provides food and cover for amphibians and reptiles, such as northern ringneck snake (*Diadophis punctatus edwardsii*) and American toad (*Anaxyrus americanus*). The eastern newt (*Notophthalmus viridescens*) spends the majority of its life as a juvenile roaming forested areas. Smaller mammals such as the gray squirrel (*Sciurus carolinensis*), eastern chipmunk (*Tamias striatus*), and raccoon (*Procyon lotor*) utilize fallen logs for cover and nest cavities. Pennsylvania's official state bird, the ruffed grouse (*Bonasa umbellus*), is an example of a non-migratory bird that prefers young, thick hardwood forest stands. Thus, they can benefit from timbering or clearing activities in the long-term when the forest regenerates back to a thick young forest stand. Upland forest is also one of several cover types inhabited by wild turkey (*Meleagris gallopavo*).

### Open Land

Several mammal species use shrublands and grasslands in the Project area as foraging and nesting habitat. These species include the eastern cottontail (*Sylvilagus floridanus*), gray squirrel, red fox (*Vulpes vulpes*), several small mammal species (e.g., mice and voles), and raccoon (NatureServe 2014). Previously cleared areas that are infrequently maintained or abandoned create successional growth habitats that are used by a distinct group of wildlife. These open lands include old agricultural fields and existing utility ROWs. ROWs in particular provide corridors that are used by several species to move between habitats. Within the Project area, these species may include white-tailed deer, coyote (*Canis latrans*), and eastern cottontail.

Other animal species such as the Virginia opossum (*Didelphis virginiana*), eastern garter snake (*Thamnophis sirtalis*), and turkey likewise rely on open land habitat for food, nesting, and protection from predators (e.g., in areas where vegetation is taller or denser than understory of adjacent habitat).

#### Forested and Scrub-Shrub Wetlands

PFO and PSS wetlands are important for providing food, shelter, migratory and wintering areas, and breeding areas for wildlife species, including red-backed salamander (*Plethodon cinereus*), wood frog (*Rana sylvatica*), and eastern garter snake. Aquatic mammals common in forested and scrub-shrub wetlands in the Project area include beaver (*Castor canadensis*), muskrat (*Ondatra zibethicus*), and mink (*Mustela vison*). Each of the wildlife species listed above for upland forests could also be present within forested and scrub-shrub wetlands.

#### Emergent Wetlands

PEM wetlands provide habitat for a variety of reptile and amphibian species, such as green frog (*Rana clamitans*), bullfrog (*Rana catesbeiana*), northern dusky salamander (*Desmognathus fuscus fuscus*), red-backed salamander, leopard frog (*Rana pipiens*), spring peeper (*Pseudacris crucifer*), northern water snake (*Nerodia sipedon*), painted turtle (*Chrysemys picta*), and snapping turtle (*Chelydra serpentina*).

#### Agricultural Areas

Although crop fields and livestock pastures do not generally provide the highest quality habitat, they often provide forage and nesting for several small mammal species such as the meadow vole (*Microtus pennsylvanicus*) that also utilize adjacent forest, open land, and even developed habitats. Additionally, pastures provide grazing habitat for species such as the whitetailed deer.

### **D3. Fishing**

**Seventy-~~Thirty~~** eight watercourses within the Project area ~~has~~ **have** a State Water Quality Designated Use Classification as Cold Water Fisheries, ~~three-ten are classified as High-~~ **Quality Cold Water Fisheries, one is are** classified as Warm Water Fisheries, and ~~six-two~~ are classified as Trout Stocked Fisheries. **South Branch Tunkhannock Creek and Tunkhannock Creek are classified by the PA Fish and Boat Commission (PFBC) as approved trout waters. The main stem of the South Branch of Tunkhannock Creek has been confirmed**

by PFBC to be stocked within 0.5 mile of the Project area. Additionally, Leonard Creek, Marsh Creek, Mill Creek, Mill Run, Whitelock Creek and their tributaries is are classified by PFBC under PA Code Title 58, Chapter 57.11 as a Wild Trout Stream, or a stream that supports the natural reproduction of trout. Additionally, three UNTs to Leonard Creek are designated by PFBC as Class A Wild Trout Stream within the Project area. South Branch Tunkhannock Creek and Martin Creek are classified by the PA Fish and Boat Commission (PFBC) as approved trout waters. The main stem of the South Branch of Tunkhannock Creek has been confirmed by PFBC to be stocked within 0.5 miles of the Project area. Additionally, Mill Creek and Whitelock Creek are classified by PFBC under PA Code Title 58, Chapter 57.11 as a Wild Trout Stream, or a stream that supports the natural reproduction of trout. The Susquehanna River and Tunkhannock Creek are is also listed by PFBC as a WWCW. Coldwater stream sections in the Project area support trout populations. Within the workspace of the Project in Wyoming County, 49-30 streams have been identified as having perennial flow and could, therefore, sustain populations of fish.

**Table L(c)-2  
 Representative Fish Species in Waterbodies Crossed by the  
 Atlantic Sunrise Project in Pennsylvania**

Common Name	Scientific Name
<b>Warmwater Fish</b>	
Largemouth bass	<i>Micropterus salmoides</i>
Smallmouth bass	<i>Micropterus dolomieu</i>
Rock bass	<i>Ambloplites rupestris</i>
Channel catfish	<i>Ictalurus punctatus</i>
Muskellunge	<i>Esox masquinongy</i>
Chain pickerel	<i>Esox niger</i>
Black crappie	<i>Pomoxis nigromaculatus</i>
Bluegill	<i>Lepomis macrochirus</i>
Pumpkinseed	<i>Lepomis gibbosus</i>
White perch	<i>Morone americana</i>
Yellow perch	<i>Perca flavescens</i>
Walleye	<i>Sander vitreus</i>
<b>Coldwater Fish</b>	
Brown trout	<i>Salmo trutta</i>
Rainbow trout	<i>Oncorhynchus mykiss</i>
Brook trout	<i>Salvelinus fontinalis</i>
Longnose dace	<i>Rhinichthys cataractae</i>
Eastern blacknose dace	<i>Rhinichthys atratulus</i>

**Table L(c)-2  
 Representative Fish Species in Waterbodies Crossed by the  
 Atlantic Sunrise Project in Pennsylvania**

Common Name	Scientific Name
Mottled sculpin	<i>Cottus bairdi</i>
Slimy sculpin	<i>Cottus cognatus</i>
Migratory Fish	
Striped bass	<i>Morone saxatilis</i>
Blueback herring	<i>Alosa aestivalis</i>
Alewife	<i>Alosa pseudoharengus</i>
American shad	<i>Alosa sapidissima</i>
American eel	<i>Anguilla rostrata</i>
Sources: PFBC	

**D4 and D5. Hiking and Observation (plant/wildlife)**

The proposed Project route does not cross any federal, state, or local parks or recreational facilities in Wyoming County.

**D6. Other**

The Susquehanna River, Tunkhannock Creek and South Branch Tunkhannock Creek have been identified as PFBC navigable waters within Wyoming County. No other functions, values, and uses have been identified for the streams and wetlands affected by this Project.

**E. Upstream and Downstream Property**

Adjoining property owners are provided in Section A, Attachment A-2.

**F. Other Environmental Factors Determined by Site Investigation**

The Project traverses multiple land use types but primarily includes open agricultural land and forested areas. During the Project siting process, Transco located all proposed facilities in such a manner as to minimize impacts to streams, wetlands, forested areas, public, and private lands. Where possible, Transco placed the proposed pipeline ROW within open areas, limiting stream and wetland crossings. More details related to the routing process for the Project is included in Attachment P.

**G. References**

- Commonwealth of Pennsylvania, Department of Conservation and Natural Resources. 2013. Bureau of Topographic and Geologic Survey. Available online at [www.dcnr.state.pa.us/topogeo/](http://www.dcnr.state.pa.us/topogeo/). Accessed March 2014.
- Cowardin LM, Carter V, Golet FC, LaRoe ET. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish & Wildlife Service Pub. FWS/OBS-79/31, Washington, DC.
- Fike, J. 1999. *Terrestrial and Palustrine Plant Communities of Pennsylvania*. Pennsylvania Natural Diversity Inventory.
- Pennsylvania Fish and Boat Commission (PFBC). 2014a. *Class A Wild Trout Waters*. Available from: <http://www.fish.state.pa.us/classa.pdf>. Accessed February 2015 and reviewed with PFBC.
- \_\_\_\_\_. 2014b. *Regional Listings of Approved Trout Waters*. Available from: <http://www.fish.state.pa.us/fishpub/summary/troutwaters.html>. Accessed February 2015 and reviewed with PFBC.
- \_\_\_\_\_. 2014c. *Stream Section That Supports Natural Reproduction of Trout*. March 2015. Available from: [http://www.fish.state.pa.us/trout\\_repro.htm](http://www.fish.state.pa.us/trout_repro.htm). Accessed February 2015 and reviewed with PFBC.
- Pennsylvania Natural Heritage Program (PNHP). 2015. Pennsylvania Endangered Reptile Species - bog turtle (*Glyptemys muhlenbergii*) fact sheet. Available from: <http://www.naturalheritage.state.pa.us/factsheets/11522.pdf>. Accessed June 25, 2016.
- The Pennsylvania Code. Title 25 Environmental Protection, Chapter 93. Water Quality Standards. Available from: <http://www.pacode.com/secure/data/025/chapter93/chap93toc.html/>. March 2014.

U.S. Army Corps of Engineers (USACE). 2011. Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0), ed. J. S. Wakeley, R. W. Lichvar, and C. V. Noble. ERDC/EL TR-10-1. Vicksburg, MS: U.S. Army Engineer Research and Development Center.

\_\_\_\_\_. 1987. Corps of Engineers Wetlands Delineation Manual. U.S. Army Corps of Engineers, Waterways Experiment Station, Wetlands Research Program, Technical Report Y-87-1. Vicksburg, MS. January 1987 - Final Report. 92 pp. + app.